
# MATHEMATICAL STRATEGIES MADE SIMPLE

A STEP-BY-STEP GUIDE FOR PARENTS AND FAMILIES





# LIBERTY ELEMENTARY

Lisa Rubin • Todd Shaffer

# MATHEMATICAL STRATEGIES MADE SIMPLE

# A STEP-BY-STEP GUIDE FOR PARENTS AND FAMILIES

This guidebook is dedicated to every adult who has ever said, "This is not how I learned to do math when I was in school," which is pretty much *all* of us.

Many of these strategies may be unfamiliar to you. They were unfamiliar to us when we first saw them as well.

Today's students must be able to do more than just find the answer. Anyone with a smart phone can find the answer in a few seconds.

Today's students must deeply understand number concepts and number relationships, working flexibly with mathematics to solve complex problems. Having taught children traditional algorithms in the past, and having since taught students these strategies, we believe in these methods as a means of helping all students truly understand the mathematics behind the work they are doing.

Bottom line: We want our students to love math as much as we do. We want our students to deeply understand math and be able to explain it to others.

> We no longer want to teach math how we were taught. We want to teach it even better.

Thank you for taking time to explore this book. It offers brief, but detailed explanations and illustrations of math strategies for addition, subtraction, multiplication and division. We sincerely hope it will be a valuable resource for you throughout your child's elementary school career.

# Lisa Rubin Todd Shaffer

# **Liberty Elementary**

Addition	1 Strategie	es 🕂
Addition Introduction	Page 1	
Base Ten Blocks Without Regrouping	Page 2	hundreds     tens     ones       Image: Ima
Base Ten Blocks With Regrouping	Page 3	hundreds     tens     ones       55     68
Number Line Addition	Pages 4-5	+10 +4 +4 +4 +15 +25 +29 +10 +12 +10 +12 +12 +12 +12 +12 +12 +12 +12 +12 +12
Number Line Addition Using Friendly Numbers	Page 6	+10 $+5$ $+20$ $+20$ $+5$ $+20$
Expanded Form Addition	Page 7	<b>56 + 24</b> 56 = 50 + 6 24 = 20 + 4 70 + 10 = <b>80</b>
Standard Algorithm Without Regrouping	Page 8	362 + <u>235</u> 597
Standard Algorithm With Regrouping	Page 9	$     \begin{array}{r}       1 & 1 \\       6 & 6 & 5 \\       + 2 & 6 & 9 \\       \hline       9 & 3 & 4     \end{array} $

Subt	raction Strategies	<b>—</b>
Subtraction Introduction	Page 10	C
Base Ten Blocks Without Renaming	Page 11	Nundreds     Lons     Ones       Image: Construction of the second seco
Base Ten Blocks With Renaming	Page 12	hundreds     tens     ones       Image: Ima
Number Lines to Count Back	Pages 13-14	-2 -2 58 60 62 82
Number Lines to Count On	Page 15	+20 79 99100 125
Standard Algorithm Without Renaming	Page 16	768 <u>-265</u> 503
Standard Algorithm With Renaming	Page 17	$     \begin{array}{r} 5 \\       7 \\       7 \\       7 \\       7 \\       9 \\       - 2 \\       6 \\       9 \\       4 \\       7 \\       0 \\     \end{array} $

Multiplication Strategies 🗙			
Multiplication Introduction	Page 18		
Skip Counting	Page 19	6, 12, 18, 24, 30, 36, <mark>42</mark>	
Repeated Addition	Page 20	15 x 4 = 15 + 15 + 15 + 15	
Models of Sets	Page 21		
Arrays	Page 22	3 x 6 = 18 or 6 x 3 = 18	
Number Line Multiplication	Page 23	$\begin{array}{c} +25 \\ 0 \\ 25 \\ 50 \\ 75 \\ 100 \\ 125 \\ 150 \end{array}$	
Area Models	Pages 24-25	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
Partial Products	Page 26	$35 \\ \times 14 \\ 20 (4 \times 5) \\ 120 (4 \times 30) \\ 50 (10 \times 5) \\ + 300 (10 \times 30) \\ 490$	
Standard Algorithm Without Regrouping	Page 27	4 1 2 <u>× 3</u> 1,2 3 6	
Standard Algorithm With Regrouping	Page 28	3 2 1 7 4 <u>× 5</u> 8 7 0	
Standard Algorithm With 2-digit Numbers	Page 29	1 52 <u>×48</u> 416 + <u>2080</u> 2496	

Division Introduction	<b>Division Strateg</b> Page 30	ies 🕂
Models of Sets	Pages 31-32	$( \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\$
Number Lines	Page 33	0 1 2 3 4 5 6 7 8 9 10 11 12
Arrays	Pages 34-35	28 ÷ 4 = 7 or 28 ÷ 7 = 4
Partial Quotients	Pages 36-37	$\begin{array}{cccc} 4 & 1 & 5 & 2 & 30 \\  & -1 & 2 & 0 & \frac{4 \times 30}{32} \\  & -3 & 2 & \frac{4 \times 8}{38} \\ \end{array}$
Multiplying Up	Pages 38-39	$240 \div 16$ $16 \times 10 = 160 \qquad 160$ $16 \times 5 = 80 \qquad +80$ $240$ $10 + 10 + 5 = 25$
Area Model	Page 40	$10   10   10$ $12 \times 10 =   12 \times 10 =   120$

# **ADDITION INTRODUCTION**

Learning addition is truly about much more than finding the answer. By using strategies like the ones in this book, your child will learn about concepts such as the relative values of numbers, how to decompose numbers into smaller parts, the values of digits within numbers, the structure of the place value system, and why regrouping works. This learning makes students more confident, fluent and flexible in their mathematical reasoning.

Structures of addition problems

Most addition problems simply involve combining two quantities (parts) together to find the **total, or sum**.

Amy read 54 pages yesterday and 39 more today. How many pages did she read in all?

We add:	Part	54
	+ Part	+ 39
	= Total	93

54 + 39 = 93 pages total

In some cases, problems may sound like subtraction, but the total is what is missing.

Gerald used 11 of his stamps mailing birthday invitations. He has 14 stamps left. How many stamps did he begin with?

We know: The part taken away = 11	So, we <u>add</u>	Part	11
The part left = 14 →	the parts	+ Part	+ 14
What we don't know:	together.	= Total	25
<u>the total</u> .			

11 + 14 = 25, so 25 - 11 = 14. Gerald began with **25** stamps total.

The following pages provide examples and step-by-step instructions for how to perform **5** different addition strategies your child will likely use in elementary school to help build his or her number sense and mathematical flexibility *while* solving addition problems. The traditional standard algorithm is also included, which is taught after students have a strong understanding of addition and the underlying numeracy concepts.

# **Base Ten Blocks Without Regrouping**

Students build or draw base ten models (blue blocks below) to represent the numbers being added (**addends**), and then combine the quantities to find the total, called the **sum**. Students start with no regrouping, where the total ones will not create a group of 10, or the total tens will not create a group of 100.




Solve: 24 + 15


**In Context:** Billy scored 24 points on Friday and another 15 points on Saturday in his basketball tournament. How many points did he score in both games combined?

### Steps:

- 1. Show 24 by building or drawing 2 tens (20) in the tens column and 4 ones (4) in the ones column.
- 2. Show 15 by building or drawing 1 ten (10) in the tens column and 5 ones (5) in the ones column.
- 3. Add the ones together: 4 + 5 = 9. Add the tens together: 20 + 10 = 30.
- 4. Add the tens and ones together: 30 + 9 = 39. Answer: 39



Here is how a student might represent it:



### Tip:

Students may be observed adding the tens first, before adding the ones. This can yield the correct answer, but remind students to watch for possible regrouping, where 10 ones will create a ten, or where 10 tens will create a group of one hundred. As students move closer to the traditional addition algorithm, they will want to get in a habit of working right to left, adding the ones first, then the tens, and so on.

# **Base Ten Block Addition With Regrouping**

Students build or draw base ten models (blue and gold blocks below) to represent the numbers to be added, then combine the quantities find the sum. Once students are proficient at solving problems without regrouping, they solve problems where the total ones are enough to create a group of ten, and/or the total tens are enough to create a group of one hundred. Students show these exchanges by grouping 10 ones to make a ten rod, or 10 ten rods to make a one hundred block (called a "flat").



Solve: 55 + 68


**In Context:** Jan walked 55 laps at Relay for Life. Sara walked 68 laps. How many total laps did the girls walk for their team?

Steps:	
oleps.	

- 1. Show 55 by building or drawing 5 tens (50) in the tens column and 5 ones (5) in the ones column.
- 2. Show 68 by building or drawing 6 tens (60) in the tens column and 8 ones (8) in the ones column.

hundreds	tens	ones	
			55
			68 ↓

- 3. Add the ones together. 5 + 8 = 13
  4. Circle 10 of the ones blocks to form an additional tens rod in the tens column. After creating a new ten, cross out the 10 ones that were circled.
- 5. Add the tens together. 50 + 60 + 10 = 120
- Circle 10 of the tens rods to form an additional hundreds flat in the hundreds column. After creating a new hundred, cross out the 10 tens that were circled.
- 7. Add 100 + 20 + 3 = **123**. **55 + 68 = 123**.




# **Adding Using Number Lines**

Students draw number lines. The numbers to be added are **decomposed** (broken apart) and shown as jumps along the line. Students can start out at one of the two numbers to be added, start out at a part of one of the numbers, or start out at 0. Once all parts of the numbers have been added, the stopping point is the total.



**Solve:** 15 + 14


**In Context:** Brittany earned 15 points on a video game on Monday. On Tuesday, she earned 14 points. How many points did she earn in two days?



Turn to the following page to see the above problem solved several more ways using number lines.

Grades 1-3

is acceptable.



# **Number Lines With Friendly Numbers**


Students draw a number line. The number to be added is broken apart and represented as jumps along the line. Students can make steps easier by adding friendly numbers (using jumps designed by the student to land on round numbers, such as 47 + 3 = 50), or by chunking numbers into any smaller parts that are easier to add quickly and efficiently using mental math.



### Example #1:

### Solve: 35 + 8

**In Context:** Juan and Danny have a dog walking business. Juan earned \$35. Danny worked less, so he earned just \$8. How much money did they earn together?



- Students are encouraged to draw larger "jumps" to show larger amounts such as +10, and draw smaller "jumps" for smaller amounts such as +2, but the sizes of jumps do not need to be perfectly proportional.
- Students may use friendlier numbers and/or larger jumps, based on their own readiness.

# Using Expanded Form to break larger numbers into smaller round<br/>numbers that are easier to mentally add or subtract. This also helps to develop<br/>mental math skills.The expanded form of a number is the value of each digit written out separately<br/>in an addition expression. Example: The expanded form of 347 is 300 + 40 + 7.Image: Colspan="2">Image: Colspan="2">Solve: 65 + 27In Context: At 9:00 AM, the temperature was 65 degrees Fahrenheit. By 2:00 PM, the temperature<br/>had increased by 27 degrees. What was the temperature at 2:00 PM?Image: Colspan="2">G5 + 27Steps:<br/>1. Find the expanded forms of 65 and 27.<br/>65 can be broken into 60 + 5Image: Colspan="2">Colspan="2">Colspan="2">Colspan= 65 + 271. Find the expanded forms of 65 and 27.<br/>65 can be broken into 60 + 5Image: Colspan="2">Colspan= 65 + 271. Find the expanded forms of 65 and 27.<br/>65 can be broken into 60 + 5Image: Colspan="2">Colspan= 65 + 271. Find the expanded forms of 65 and 27.<br/>65 can be broken into 60 + 5Image: Colspan="2">Colspan= 65 + 271. Find the expanded forms of 65 and 27.<br/>65 can be broken into 60 + 5Image: Colspan="2">Colspan= 65 + 272. Steps:<br/>0. Steps:<br/>0. Steps:Image: Colspan="2">Colspan= 65 + 271. Find the expanded forms of 65 and 27.<br/>0. Steps:<br/>0. Steps:Image: Colspan="2">Colspan= 65 + 272. Steps:<br/>0. Steps:I

- 2. Add the tens from both numbers.<br/>60 + 20 = 8065 = 60 + 5<br/>7 = 1227 = 20 + 7<br/>Tens  $\rightarrow$ 3. Add the ones from both numbers.<br/>5 + 7 = 1260 + 20 = 80<br/>Ones  $\rightarrow$ <br/>5 + 7 = 12
- Add the total tens and ones together to find the sum.
   Answer: 80 + 12 = 92.

27 can be broken into 20 + 7

There is not one "right way" to write down the steps taken in this method. Students can organize the steps however they like.

**80 + 12** = 92

Here, the expanded forms of the numbers are stacked in columns for easy addition.	In this example, the expanded forms are organized with the hun- dreds first, then the tens, and then the ones in an expression.	
56 + 24	264 + 315 =	
56 = 50 + 6	(hundreds) (tens ) (ones)	
24 = 20 + 4	200 + 300 + 60 + 10 + 4 + 5 =	
70+10=80	500 + 70 + 9 =	
70 + 10 - 80	579	

**Tip:** As with number lines, remember that this approach is not about solving efficiently at first. It is about helping students understand place value and increasing mathematical flexibility.

# Standard Algorithm Without Regrouping

The standard algorithm is used when students have a deep understanding of place value. These examples have no **regrouping** (or what many adults called "carrying the one" when we were growing up in school).



### Solve: 123 + 245

**In Context**: A third grade class donated 123 books to a book sale. A fourth grade class donated 245 books to the book sale. How many books were donated to the book sale?

### Steps:

1.	. Stack the two numbers neatly above one another.	123
	Line up digits in the ones, tens, and	+245
	hundreds places under the corresponding place values.	

2.	Add the digits in the ones place. $3 + 5 = 8$	123
3.	Add the digits in the tens place. $2 + 4 = 6$ (20 + 40 = 60)	+245
4.	Add the digits in the hundreds place. $1 + 2 = 3$ (100 + 200 = 300)	368

### Answer: 368

Tips:

- Remind children of the values of the digits. For instance, after solving the above example,
   Ask: What is the value of the 6 in your answer? (60)
   How did we get 60? (adding 20 + 40)
- Sometimes a student may incorrectly stack the numbers, such as in this example:

243 +25

This may reveal that a child does not yet have a clear and complete understanding of place values. If this occurs, it may be time to revisit expanded form, base 10 models, or other concrete representations.

# **Standard Algorithm With Regrouping**

The standard algorithm is used when students have a deep understanding of place value. While it does not show the true value of the digits, it can be an efficient strategy for students who have a solid background. Once regrouping is involved, it becomes even more crucial that students know that ten ones make 10, ten tens make 100, etc.




### **Solve**: \$769 + \$165

**In Context**: Malcolm saved his money for several months. When he went to an auction, he was able to buy the go-cart he had dreamed of for \$769. He still had \$165 left. How much money had he saved <u>before</u> going to the auction?

Steps:	Steps:				
up t	ck the two numbers neatly above one another. Take care to line he digits in the ones, tens, and hundreds places under the responding place values.	769 +165			
3. Don mak	the digits in the ones place. 9 + 5 = <b>14 ones</b> It write 14 in the ones place below the line. Regroup 10 ones to a ten. Write the 1 above the tens column (over the 6 as shown in Then write the remaining 4 ones below the line in the ones place.	1 769 +165 4			
5. Don a hu sho	the digits in the tens place. 6 + 6 + 1 = <b>13 tens</b> "'t write 13 in the tens place below the line. Regroup 10 tens to make undred. Write the 1 above the hundreds column (over the 7 as wn in green). Then write the remaining 3 tens below the line in the s place.	1 1 769 +165 34			
Wri	the digits in the hundreds place. 1 + 7 + 1 = <mark>9 hundreds</mark> . te the 9 hundreds below the line in the hundreds place. swer: 934	<sup>1 1</sup> 769 +165 934			

Tip:

Remind your child to only regroup when there is at least ten in any given place value. Some problems may involve regrouping only the tens, only the hundreds, or neither.



# **Base Ten Blocks Without Renaming**

Students build or draw models to represent subtraction using a "take from" approach using blocks or pictures of base ten blocks. Students start with no renaming ("borrowing"). The starting number (**minuend**) has enough ones and tens to give away during the subtraction. Students take away or cross out the amount to be subtracted in order to find the amount left, called the **difference**.



ones

 $\sim$ 

 $\square$ 

 $\square$ 

Solve: 68 — 42 minuend subtrahend

hundreds

**In Context:** Marta has gotten an impressive 68 base hits this season. She had 42 hits for her team last year. How many more hits has she gotten this year compared to last year?

### Steps:

 Build or draw the starting number (minuend). Show 68 by building or drawing 6 tens (60) in the tens column and building or drawing 8 ones (8) in the ones column.

Students do not need to build or draw the 42.

- Take away the 42 (subtrahend).
   Remove/cross out 2 ones (2).
   Remove/cross out 4 tens (40).
- Count the number of tens that are left.
   (2 tens = 20)
- 4. Count the number of ones that are left.(6 ones = 6)

Answer = 68 - 42 = 26

hundreds	tens	ones
	<i>11111</i> 11	

tens

Tip:

• Students may be observed subtracting the tens first, before subtracting the ones. This can yield the correct answer when no renaming is needed, but not when there are not enough of a given place to take away. Students will want to quickly form a habit of working right to left, subtracting the ones first, then the tens, etc.

Grades 1-3

# **Base Ten Blocks With Renaming**

Students build or draw models to represent subtraction using a "take from" approach using blocks or pictures of base ten blocks. Here, students solve problems where the number of ones or tens in the starting number (minuend) is less than the ones or tens in the number to be taken away (subtrahend). Students must "borrow" by replacing a ten rod to make 10 ones, or a 100 block to make 10 tens, and so on. Students then take away or cross out the amount to be taken away (subtrahend) in order to find the amount left (difference).



hundreds

**In Context:** Mary flipped a coin 45 times. The coin came up heads 27 times. How many times did her coin show tails?

### Steps:

- Build or draw the starting number (minuend). Show 45 by building or drawing 4 tens (40) in the tens column and building or drawing 5 ones (5) in the ones column. Students do not need to build the 27.
- hundreds
   tens
   ones

   Image: Constraint of the second second

tens

ones

- 2. There are not enough ones in 45 to remove or cross out 7 ones.
- 3. Rename a ten. Move 1 ten (10) from the tens column and replace it with 10 ones (10) in the ones column.


Notice that there is still a value of 45 cubes shown at this point. 3 tens (30) and 15 ones (15) is still 45.

4. Now there are 15 ones, so you can subtract the 7 ones in 27.

Begin to take away the 27 (subtrahend).

Remove/cross out 7 ones (7).

Remove/cross out 2 tens (20).



5. Count the number of tens that are left (1 ten = 10).

6. Count the number of ones that are left (8 ones = 8).

Answer = 45 - 27 = 18

Tip: Students must work right to left, subtracting the ones first, then the tens, etc.

Grades 2-3

# **Number Lines to Count Back**

Students draw a number line and then represent subtraction of the decomposed parts of the numbers as **backwards** jumps along the line. Once all parts of the numbers have been subtracted, the stopping point is the answer (called the difference).



	<b>Solve:</b> 82 - 24				
In	In Context: There were 82 students in the fourth grade. 24 of the students did their homework for				
ex	tra credit. How many students will not receive extra credit?				
He	ere is one way to solve this problem with a number line.				
Ste	eps:				
1.	Draw a blank number line and place the starting number (82) on the right, leaving space to the left to count backwards.				
2.	We need to jump backward 24. There are many ways to decompose the 24. One way is to subtract 24 by going backwards 2 jumps of 10 and then 4 jumps of 1.				
3.	Draw and label a backwards jump of 10 on the number line. Land on 72.				
4.	Draw and label 2nd backwards jump of 10 on 62 72 82 the number line. Land on 62.				
5.	Draw and label four backwards jumps of 1 along the number line. $-10 -10$				
	We go back to 61, 60, 59, and finally 58. Answer: 58				
Th	In the above example, we subtracted 82 - 24 by starting at 82 and subtracting 2 tens and 4 ones. There are many other correct ways to solve this problem on a number line. As with addition, there is typically always more than one path to the solution, and any efficient and accurate path is				

Turn to the following page to see the above problem solved two more ways using number lines.

acceptable.

# Number Lines to Count Back (continued)


Solve: 82 - 24

**In Context:** There were 82 students in the fourth grade. 24 of the students did a project for extra credit. How many students will not receive extra credit?




Here is the same problem, 82 - 24, solved two more correct ways:

On this number line, the student decomposed the 24 into 10, 10, 2, and 2. This takes advantage of friendly numbers. By splitting the 4 into 2 and 2, the movement backwards across 60 is easier.



On this number line, the student moved backwards by 20 in one jump. Then, the student used friendly numbers, splitting up the 4 into 2 and 2. The student subtracted 2 to land on 60, before going back 2 more to the answer of 58.



Tips:

- Encourage students who may be subtracting with only jumps of one to begin taking larger jumps as they show they are ready. As students gain experience and confidence, they can combine numbers with more efficient and clever combinations.
- As with addition, students can mentally decompose numbers before and during subtraction, in
  order to use a more strategic approach. Students can make steps easier by subtracting friendly
  numbers, such as 10 or 20. Students can also use jumps they design to land on round numbers,
  such as changing 43 6 into 43 3 3, so as to land neatly on 40 before going back further.

# **Number Lines to Count On**

# (Finding the Difference)

Not all subtraction involves "taking away from." Many subtraction problems involve finding a *difference* between two numbers: 85 - 83 = 2 *because 85 and 83 are 2 jumps apart*. Therefore, students can also solve subtraction problems on a number line by finding how many jumps are *between* the two numbers. To do this, students start at the smaller number (subtrahend) and draw and label jumps forward until the larger number (minuend) is reached. The total of the jumps will tell you the difference (your answer).



# Solve: 125 - 79


minuend subtrahend

**In Context:** There are now 125 new cars on the car lot. Before today's delivery arrived, there were only 79 cars on the lot. How many new cars were delivered today?

### This is one way to solve this problem with a number line:

### Steps:

- 1. Draw a number line and mark 79 as a starting point, leaving space to count on.
- 2. Jump ahead by tens until to get as close to the target of 125 as possible without going over.
- 3. Jump ahead by ones until you reach 125.




4. Add up the numbers that were added to get to 125.

The total difference between 79 and 125 is 46. Answer: 46

During the delivery, 46 cars were added to the original 79 to reach the current total of 125.

The same problem solved with some larger and more efficient jumps:



This student added 20 and then 1 more, in order to land on 100. Then another jump of 25 was needed, going from 100 to 125. The total difference (20 + 1 + 25) between 79 and 125 is still **46**.





# Standard Algorithm Without Renaming

Grades 4-5

The standard algorithm is used when students have a deep understanding of place value. These examples have no renaming (some people call this "borrowing").



Solve: 68-23

**In Context:** In a pet store, there were 68 animals to be adopted. 23 families adopted pets over the weekend. How many animals were still waiting to be adopted?

Steps: 1. Stack the two numbers neatly above one another. Line up digits in the ones, tens, and hundreds under the corresponding place values.	68 - 23			
Because the digits in the top number (minuend) are all larger than the correspondin bottom number (subtrahend), no renaming is needed to solve this problem.	g digits in the			
<ul> <li>2. Subtract the digits in the ones place. 8 - 3 = 5</li> <li>3. Subtract the digits in the tens place. 6 - 2 = 4 (60 - 20 = 40)</li> <li>Answer: 45</li> </ul>	68 - 23 45			
<ul> <li>Tips: <ul> <li>Remind children of the values of the digits. For instance, after solving the problem above: You might ask your child questions such as:</li> <li>What is the value of the digit 4 in your answer? (40) How did we get 40? (subtracting 60– 20)</li> </ul> </li> <li>Sometimes a student may incorrectly stack the numbers, such as in this example: <ul> <li>243</li> <li>-21</li> </ul> </li> <li>This may reveal that a child does not yet have a clear and complete understanding of place values. If this occurs, it may be time to revisit expanded form, base-10 models, or other concrete representations.</li> </ul>				

# **Standard Algorithm With Renaming**

The standard algorithm is used when students have a deep understanding of place value. While it does not show the true value of the digits, it can be an efficient strategy for students who have a solid background. Once renaming is involved, it becomes even more crucial that students know that ten ones make 10, ten tens make 100, etc.



## **Solve**: 234 — 56

**In Context:** After the farm market closed, the Anderson family had 56 jars of their world famous jams and jellies remaining. They began the weekend with 234 jars. How many jars did they sell?

1.	<ul> <li>Stack the two numbers above one another. Make sure the greater number is on the top. Line up digits in the ones, tens, and hundreds under the corresponding place values.</li> <li>Start with the ones column. Subtract the bottom number from the top number. Can you subtract 4 - 6? No. There are not enough ones to take 6 away.</li> </ul>	234 - 56
	Go to the tens column. Transfer 1 ten (worth 10 ones) from the tens column. Show this by marking out the 3 and showing that there are now only 2 tens left in the tens column. We are trading this ten for 10 ones. Add the 10 ones that were transferred to the ones column: (10 + 4 = 14)	2 <sup>14</sup> 2 <b>3</b> 4 - 56
5.	You have 14 ones so you can take 8 away. Now subtract the ones: $14 - 6 = 8$ . Look at the tens column. Subtract the bottom tens digit from the top tens digit if you can. Can you subtract 2 tens - 5 tens? No. There are not enough tens to take 5 tens away.	8
6.	Go to the hundreds column. Transfer 1 hundred (worth 10 tens) from the hundreds column. Show this by marking out the 2 and showing that there is now only 1 left in the hundreds place.	1 12 14 234 - 56
7.	Add the 10 tens that were transferred to the tens column. $10 + 2 = 12$ . Subtract the tens: 12 tens - 5 tens = 7 tens.	<u>78</u>
8.	Look at the hundreds column. Subtract the bottom hundreds digit from the top hundreds digit.	1 12 14 <b>XXX</b>
9.	Subtract 1 hundred - 0 hundreds.	- 56
	100 - 0 = 100.	178
	Answer: 178	
Gra	ades 4-5	

# **MULTIPLICATION INTRODUCTION**

Multiplication is a way to quickly find the total amount in a collection of equal groups or sets. Let's take a quick look at the 3 most common situations that involve multiplication.

### 1. Equal Groups or Sets

In many multiplication problems, there are a given number of groups or sets, each of which is equal in size or quantity. Multiply the number of groups × the number in each group in order to find the total (product).

Derrick bought 6 packs of crayons. Each pack of crayons has 8 crayons inside. How many crayons did he buy altogether?



number of groups	×	number in each group	= total
6		8	= 48
number of boxes	×	crayons in each	total
			crayons


6 packs of 8 crayons make 48 crayons in all.

6 × 8 = 48

# 2. Arrays

Arrange items into equal rows and columns to form a rectangular array. Rather than counting all of the items in the array, you can multiply the number of rows × the number of columns, in order to find the total (product).

The chairs for the meeting were arranged in 3 rows of 9. How many chairs were set up?



# **Skip Counting to Multiply**

Skip counting means counting by the same number repeatedly. Example: Skip counting by three: 3, 6, 9, 12, 15, 18, 21... This method is used to solve basic multiplication problems. Students can skip count the correct number of jumps to find the total. The total in multiplication is called the **product**.



**Solve:** 4 X 5

**In Context:** At Freedom Elementary School, there are 4 classrooms in the second grade. Each classroom has 5 class pets. How many pets are in the second grade classes?

St	eps:							
1.	Think about the problem:							
	Classroom #1 has 5 pets	Classroom #2 has	s 5 pets					
	Classroom #3 has 5 pets	Classroom #4 has	s 5 pets	i -				
2.	Skip count by fives (4 jumps) to find t				<b>5</b> , (1x5)	<b>10</b> , (2x5)	15, <sup>(3x5)</sup>	<b>20</b> (4x5)
	Answer: I	There are <b>20</b> pets	altogeth	ner.				
	Here	e is one more exa	mple:					
	6 friends each eat 3 cookies	. How many cool	kies wer	e eaten	n? <b>S</b> o	<b>olve:</b> 6 :	х З	
St	eps:							
1.	Think about the problem:							
	Friend #1 has 3 cookies Friend	#2 has 3 cookies	Fri	end #3	has 3	cookie	S	
	Friend #4 has 3 cookies Friend	#5 has 3 cookies	Fri	end #6	has 3	cookie	S	
2.	Skip count by threes (6 jumps) to find (total).	I the product	<b>3</b> , (1x3)	<b>6</b> , <sup>(2x3)</sup>	9, <sup>(3x3)</sup>	12, (4x3)	15, <sup>(5X3)</sup>	<b>18</b> (6x3)
	Answer: 18	cookies were eate	en altog	ether.				
	Note: While early learners of multiplication may not recognize this at first, this problem could also be solved in fewer jumps by counting by 6, three times.							
	6, 12, <b>1</b>	<b>8</b> . (3 x 6 =	= 6 x 3)					
Tip	<b>):</b> This method typically only applies to sc	olving basic multip	lication	math fa	acts a	nd patte	erns.	

# **Repeated Addition to Multiply**

As students are learning the concept of multiplication, they must understand that multiplication is a way to combine equal groups or sets. Repeated addition is just as it sounds: adding the same number repeatedly in order to find the total.



## Solve: 17 X 4

**In Context:** A group of students filled 4 bags of apples at the apple orchard. Each bag contained 17 apples. How many apples did the students pick?

### Steps:

1. Think about the problem:

Bag #1 contained 17 apples Bag #3 contained 17 apples Bag #2 contained 17 apples Bag #4 contained 17 apples

2. Write out the multiplication problem as repeated addition.

 $17 \times 4 = 17 + 17 + 17 + 17$ 

3. Use any addition strategies to solve the problem. Here are three different methods:

17 + 17 = 34	<sup>2</sup> 17 17	17 + 17 + 17 + 17 =
34 + 17 = 51	17	34 + 34 =
51 + 17 <b>= 68</b>	<u>+17</u> <mark>68</mark>	68

The students collected 68 apples.

### Tips:

• Third graders learn about the commutative property of multiplication. Repeated addition is another way for students to learn the importance of the commutative property of multiplication, which shows that 4 x 13 is the same as 13 x 4.

 Repeated addition helps illustrate the meaning of the multiplication operation. Understanding the repeated addition concept will also have value for students IN LATER GRADES as they first begin to learn how to multiply fractions and decimals by whole numbers in grades 4-5.


> Grade 4 Application:  $3 \times \frac{1}{4} = \frac{1}{4} + \frac{1}{4} + \frac{1}{4} = \frac{3}{4}$ Grade 5 Application:  $5 \times 0.8 = 0.8 + 0.8 + 0.8 + 0.8 + 0.8 = 4.0$

Grades 2-3

# **Using Models of Sets to Multiply**

In this early and important multiplication method, students build or draw a model of equal sets to represent and solve multiplication problems. Sets must be equal in size.

This method can be used to solve basic multiplication problems. Students count the total number of objects in the sets to find the **product**.



### **Solve:** 4 X 6

**In Context:** Tom baked mini pizzas for the annual football party. He filled 4 trays with pizzas. Each tray held 6 pizzas. How many mini pizzas did Tom bake?

### Steps:

- 1. Draw four trays (symbols are fine).
- 2. Draw 6 pizzas in each tray.
- 3. Add or count to find the total number of pizzas.

6 + 6 + 6 + 6 **= 24** 

Answer: 24

### Another example:


### **Solve:** 5 X 7

**In Context:** There are 5 bookshelves with 7 books on each shelf. How many books are there altogether?

### Steps:


- 1. Draw five shelves (symbols are fine).
- 2. Draw 7 books in/on each shelf.
- 3. Add or count to find the total number of books.

7 + 7 + 7 + 7 + 7 = 35 Answer: **35** 



### Tips:


- Sets of everyday objects can be used to model multiplication. There is no need to rely solely on drawings. Make equal groups of pennies, toys, food...or anything around the house!
- Students may use skip counting to total the objects in the groups.
- When creating drawings, students should keep them very simple. Students can represent 3 barns with 4 horses in each barn by drawing rectangles for the barns and x's for the horses.




# **Arrays for Multiplication**

An array is a visual organization of symbols, shapes or objects in equal rows and columns. It is typically rectangular. This is a critical model for multiplication, in which:

*the number of rows × the number of columns = the total objects* Students draw arrays and then use skip counting, addition, or other strategies to total up the objects in the arrays.





22

# **Multiplying Using Number Lines**

Students can use the same number line drawings they used for addition to represent multiplication as repeated addition of equal groups. Students draw a number line and represent the equal groups or sets as equal jumps along the line, starting at 0. Once all the sets or groups have been added, the stopping point is the total.



### **Solve:** 4 × 10

**In Context:** Mark had 4 presents to be wrapped. Each present required 10 inches of ribbon. How much ribbon will Tom need?

### Steps:

- 1. Draw a blank number line and mark 0 as a starting point.
- 4 x 10 can represent four tens or ten fours, and can be shown either way on a number line. In this example, it is easier and faster to count by tens. Draw 4 jumps of 10 forward on the number line: 0, 10, 20, 30, 40.

Answer: 40



$$4 \times 10 = 10 \times 4$$

+10

+10


20

+10


+10

40

30



This is a different example using a number line to calculate 6 x 25, shown as 6 jumps of 25.




Students will be able to explore which representation will be most efficient. Imagine what 25 jumps of 6 would look like!

**Tip:** This strategy is not an efficient long-term strategy, but it helps build students' understanding of the multiplication concept, and it connects multiplication to students' understanding of addition.

# **Area Model of Multiplication**

The area model for multiplication is based on the concept of **area of a rectangle**. Area of a rectangle is calculated by multiplying length times width to find the total squares within the rows and columns. The total square units inside the rectangle are the product.

Example: 4 x 6 = 24 square units





Solve: 34 X 16


**In Context:** In a hot dog eating contest, the contestants ate 34 large packs of hot dogs, with 16 hot dogs in each pack. How many hot dogs did the contestants eat in all?

<b>C</b> t			34	
	<ul><li>Steps:</li><li>1. Draw a rectangle with the length (34) and width (16) labeled as the two factors to be multiplied.</li></ul>			
2.	Decompose/break apart 34 into friendlier numbers that can be multiplied easily in your head. Students often decompose numbers into the values of each digit, so we will just split 34 into 30 and 4. Write the 30 and 4 on the top of the area model. Draw a vertical line inside the rectangle, and try to show that 30 is larger than 4.	16	34	_4
3.	Decompose/break apart 16 into friendlier numbers that can easily be multiplied in your head. We will split 16 into 10 and 6. Draw a horizontal line inside the rectangle, and try to show that 10 is larger than 6.	1 16	34 30 6	4
	Use mental math to multiply length and width and find the ard Multiply 30 × 10 and write the product, 300, inside the upper left hand section. Multiply 30 × 6 and write the product, 180, inside the lower left hand section. Multiply 4 ×10 and write the product, 40, inside the upper right hand section. Multiply 4 × 6 and write the product, 24, inside the lower right hand section. Add up the 4 smaller areas to find the total: 300 + 180 + 40		34 30 300 30 180	4 40 4 24
	des 3-5	, T <b>2</b> 4	- J Aliswei. 544	 _

# **Area Model for Multiplication (Continued)** The area model for multiplication is based on the concept of area of a rectangle. Area of a rectangle is calculated by multiplying length to width to find the total squares within the rows and columns. The total square units inside the rectangle are the product. An example with a 3-digit number multiplied by a 1-digit number: Solve: 159 × 6 Steps: 1. Draw a rectangle with the length (159) and width (6) labeled as the two factors to be multiplied. 159 6 2. Break apart 159 into friendlier numbers that can be multiplied easily in your head: 100 + 50 + 9. Write the 100, 50, and 9 on the top of the area model. Draw vertical lines inside the rectangle to separate the 3 sections. Write the other factor, 6, on the side as the width. 159 100 6 3. Mentally multiply length x width to find the area of each section. Multiply 100 × 6 and write the product, 600, inside the 1st section. Multiply $50 \times 6$ and write the product, 300, inside the 2nd section. Multiply 9 × 6 and write the product, 54, inside the 3rd section. • 159 6 54 600 300 4. Add up the 3 smaller areas to find the total: 600 + 300 + 54 = 954 Answer: 954

### Tip:

Graph paper is a great tool for creating area models. It is particularly useful when students are first learning about the area model, or when multiplying larger numbers. Trace around the area you want, in order to make a quick model that actually shows the square units. Students can also trace and label smaller sections of the figure and use partial products to solve a large area on graph paper, adding the areas of the pieces together to find the total area of the figure.



Partial Products (Usin Partial products is a strategy where students numbers into smaller, round numbers that are Then students add the partial products togeth The expanded form of a number is the value addition expression. Example: The	use <b>expanded form</b> to break larger e easier to mentally multiply together ner to find the total.	n an	
	<b>Solve:</b> 35 X 14		$\sim$
In Context: There were 14 girls on the tra 35 laps around the track in fifty minutes.	ack team, who each set a goal to	>	35 = 30 + 5 14 = 10 + 4
Steps:		35 4	
<ol> <li>Decompose/break apart 35 into 30 + 4</li> <li>Decompose/break apart 14 into 10 + 4</li> </ol>		×14	-
<ul> <li>3. Multiply each value from 35 (30 and 5 4).</li> <li>Multiply 4 (from 14) X 5 (from 35). Th</li> <li>Multiply 4 (from 14) X 30 (from 35). T</li> <li>Multiply 10 (from 14) X 5 (from 35). T</li> <li>Multiply 10 (from 14) X 30 (from 35).</li> </ul>	is equals 20. his equals 120. his equals 50.	35 ×14 20 120 50 300	(4 x 5) (4 x 30) (10 x 5) (10 x 30)
<ol> <li>Add all of the products together. 300</li> <li>Answer: 490</li> </ol>	+ 50 + 120 + 20 = <b>490</b>	35 ×14 20 120 50 +300 490	(4 x 5) (4 x 30) (10 x 5) (10 x 30)
<ul> <li>Tips:</li> <li>Students can organize the steps howe stacking makes the addition of the part</li> <li>Each digit from the bottom number mu <i>However, it may be wise to do the step standard algorithm to avoid confusion</i></li> </ul>	tial products easier. st be multiplied by each digit in th os <i>in the same order as you would</i>	e top numb	er.

Grades 4-5

# Standard Algorithm Without Regrouping

The standard algorithm is used when students have a deep understanding of place value, multiplication, and how the digits represent equal groups. This strategy is often confusing for students who don't yet have these concepts.



Solve: 412 X 3

In Context: The farmer's new front fence will be 412 yards long. A yard is equal to 3 feet.<br/>How many feet long will the front fence be?412<br/> $\times 3$ Steps:<br/>1. Stack the two numbers neatly above one another. Take care to line up digits<br/>in the ones, tens, and hundreds under the corresponding place values.412<br/> $\times 3$ *Multiply 3 x 412 by multiplying 3 times each digit in the top number.*412<br/> $\times 3$ 2. Multiply the digit 3 in the ones place in the bottom number by the digit 2 in412<br/> $\times 412$ 

2.	the ones place in the top number. $3 \times 2 = 6$ (ones)	$\times 3$
3.	Write the product, 6, in the ones place underneath the ones column.	6
4.	Multiply the digit 3 in the ones place in the bottom number by the digit 1 in the tens place in the top number. $3 \times 1 = 3$ (tens)	412 × 3
5.	Write the product, 3, in the tens place.	<mark>3</mark> 6
	Multiply the digit 3 in the ones place in the bottom number by the digit 4 in the hundreds place in the top number. $3 \times 4 = 12$ (hundreds)	412 × 3
7.	Write the product, 12, starting in the hundreds place. Because you have 12	× 3

hundred, the answer spills over into the thousands place. So, the 2 hundreds **1,236** go beneath the hundreds column, and the 1 thousand goes to the left.

Answer: 1,236


**Tip:** If your child becomes confused by the steps, try solving the problem using the partial products method on the previous page. Then retry the problem with the standard algorithm, making connections between the steps.

412	<b>412</b>
× 3	<u>× 3</u>
6 (3 x 2)	1,236
30 (3 x 10)	$3 \times 2 = 6$
6 (3 x 2) 30 (3 x 10) 1200 (3 x 400)	$3 \times 2 = 6$ $3 \times 10 = 30$ $3 \times 400 = 1200$
1,236	$3 \times 400 = 1200$

27

# Standard Algorithm With Regrouping

The standard algorithm is used when students have a deep understanding of place value, multiplication, and how the digits represent equal groups. While it does not show the true value of the digits, it is an efficient strategy for students who have a solid background.




### **Solve:** 174 X 5

**In Context:** The delivery truck is loaded up with 5 new refrigerators. These lightweight refrigerators each weigh 174 pounds. What is the total weight of the cargo?

	eps: Stack the two numbers neatly above one another. Line up digits in the ones, tens, and hundreds under the corresponding place values.	174 × 5
	Multiply 5 x 174 by multiplying 5 by each digit in the top number.	
2.	Multiply the digit 5 in the ones place in the bottom number by the digit 4 in the ones place in the top number. $5 \times 4 = 20$ ones	2 174
3.	Don't write 20 in the ones place below the line. Regroup the 20 ones to make 2 tens and 0 ones. Write the 2 above the tens column (over the 7 as shown in red). Write the remaining 0 ones below the line in the ones place.	× 5 0
4.	Multiply the digit 5 in the ones place in the bottom number by the digit 7 in the tens place in the top number. $5 \times 7 = 35$ tens Add the 2 extra tens on top from the previous step. $35 + 2 = 37$ tens	<mark>32</mark> 174
5.	You cannot write 37 in the tens place below the line. Regroup the 37 tens (370) to make 3 hundreds and 7 tens. Write the 3 above the hundreds column (over the 1 as shown in blue). Write the remaining 7 tens below the line in the tens place.	× 5 70
6.	Multiply the digit 5 in the ones place in the bottom number by the digit 1 in the hundreds place in the top number. $5 \times 1 = 5$ hundreds Add the 3 extra hundreds on top from the previous step. $5 + 3 = 8$ hundreds	32 174 × 5
7.	Write the 8 hundreds in the hundreds place.	870
	Answer: 870	
Tip	p: Remember to only regroup to the top of the next place value when there is at least ten in any given place value at the end of a step. Some problems may involve regrouping only the tens, or only the hundreds, or neither.	

# **Standard Algorithm With 2-Digit Numbers**


The standard algorithm is used when students have a deep understanding of place value, multiplication and how the digits represent equal groups. While it does not show the true value of the digits, it can be an efficient strategy for students who have a solid background.



### **Solve:** 52 x 48

**In Context:** José and Marcus are attempting to build a record size house of cards. They hope to use 48 decks of cards. Each standard deck contains 52 cards. How many cards will be in the card house if they are successful?

Steps: 52			
1.	Stack the two numbers neatly above one another. Line up digits in the ones, tens, and hundreds under the corresponding place values.	× 48	
	First, multiply 8 decks of cards x 52 cards in each.		
	Multiply the digit 8 in the ones place in the bottom number by the digit 2 in the ones place in the top number. $8 \times 2 = 16$ ones Don't write 16 in the ones place below the line. Regroup the 16 ones to make 1 ten and 6 ones. Write the 1 above the tens column (over the 5 as shown in red). Write the remaining 6 ones below the line in the ones place.	1 52 × 48 6	
Tip	Multiply the digit 8 in the ones place in the bottom number by the digit 5 in the tens place in the top number. $8 \times 5 = 40$ tens Add the 1 extra ten on top from the previous step. $40 + 1 = 41$ tens cross out the 1 after you add it, so you don't accidentally add it again in the next step. Write the 41, starting in the tens place. Because you have 41 tens, this spills over into the hundreds place.	1 52 × 48 416	
	Next, multiply the other 40 decks of cards x 52 cards in each.		
	Multiply 40 by 52. Start by placing a 0 underneath the 6 in the ones place of the answer section beneath the line. This 0 is called a placeholder. It represents the 0 in 40, because you are multiplying by 52 by 40, not just 4. Multiply the digit 4 in the tens place in the bottom number by the digit 2 in the ones place in the top number. $4 \times 2 = 8$ tens. Write the 8 in the tens place next to the placeholder 0.	× 52 × 48 416 80	
9. 10 11	Multiply the digit 4 in the tens place in the bottom number by the digit 5 in the tens place in the top number. $4 \times 5 = 20$ hundreds. Don't add the old 1 on top! Write the 20, starting in the hundreds place. Because you have 20 hundreds, this spills over into the thousands place. Add the 416 (from 8 x 52) plus the 2,080 (from 40 x 52). $(8 \times 52) \longrightarrow + (40 \times 52) \longrightarrow + (40 \times 52)$	× 48 × 48 416 2080 2,496	



# **Using Models of Sets to Divide**

In this early division model, students build or draw models of equal sets to represent and solve division problems. They may draw equal groups and distribute all of the items into each group, or make all the groups of a given size, until they reach the total. The answer is called the **quotient**.




### **Example #1:** Solve: 18 ÷ 3

**In Context:** There are 18 students in a science classroom. They are put into three equal groups for a class project. How many students will be in each group?

### Steps:

1. The problem states that there are 3 groups, and we need to find how many are in each group.



- 2. To solve this problem, start by drawing 3 shapes, such as circles or boxes, to represent each of the 3 groups.
- 3. Add one student into each group. Represent 18 students with stars or other symbols until all 18 students are placed in groups.
- 4. Count the number of students (stars) in each group.The quotient (answer) is 6. There are 6 students in each group.

### **Example #2:** Solve: 40 ÷ 5

**In Context:** Turtletown Elementary needs to order spiral notebooks for all 40 of its 3rd graders. The notebooks come in packs of 5. How many packs of notebooks will they need to order so that all 40 students receive one?

### Steps:

1. This problem states that there are equal groups of 5. We need to find how many groups of 5 make 40.

Number?f Groups × Number in Each Group

- 2. Draw a quick shape, such as a circle or box, to represent a pack of 5.
- 3. Draw another pack, and then another, keeping track of how many total notebooks are included, until there are 40 total notebooks.
- 4. Count how many packs were made. The quotient (answer) is 8. It takes 8 packs of 5 to make a total of 40 notebooks.

 $\Delta \Delta \Delta \Delta \Delta$ 

Grade 3

# Using Models of Sets to Divide (Continued with remainders)

Some problems have numbers that may not divide evenly, such as 11 divided by 5. After forming equal groups, a partial group of extra items remains. The number of leftover items that cannot form a complete group or cannot be shared with each existing group is called the **remainder**.

Solve: 20 ÷ 3

Example #3 with a remainder:

**In Context:** There are 20 students in a science classroom. They are put into three equal groups for a class project. How many students will be in each group?

### Steps:

- 1. The problem states that there are three groups, but we want to find out how many are in each group.
- 2. Draw some quick shapes, such as a circles or boxes, to represent each of the 3 groups.
- 3. Add one student into each group (students are represented with stars) until all 20 are placed.
- Count the number of students (stars) in each group.
   The number 20 does not evenly split into three groups, so you can put 6 students in each group, but this problem has a remainder of 2 extra students.
- 5. The quotient (answer) is 6 with a remainder of 2. There are 6 students in each of the 3 groups, with two students left over. (Students may realize that two of the groups would have to have 7 students.)

**Solve:** 43 ÷ 5

Example #4 with a remainder:


**In Context:** Turtletown Elementary needs to order spiral notebooks for all 43 of its 3rd graders. The notebooks come in packs of 5. How many packs of notebooks will they need to order?

### Steps:

- 1. This problem states that there are equal groups of 5, but we want to find out how many groups of 5 will make 43.
- 2. Draw quick shapes, such as boxes, to stand for the packs of 5. Draw another pack, and then another, until you reach 43 total notebooks.
- It takes 8 packs of 5 to make a total of 40 notebooks, but 9 packs would be 45. It is impossible to make exactly 43 using packs of 5.

The answer to  $43 \div 5$  is <u>8 with a remainder of 3</u>. The answer to the *question* of how many packs to order is that the school would need to order <u>9 packs</u>, so everyone gets a folder.






You would need to buy an extra pack to get the 3 extra notebooks you need.





### This strategy provides students with a visual model of division. Example #1 Solve: \$12 ÷ 4 In Context: John has \$12. He wants to buy some books that each cost \$4. How many books can he buy? Steps: 1. Draw a number line and label it with marks for every number from 0 to the total, which is 12. Ś 5 6 8 9 10 11 12 2. Draw as many jumps of 4 as possible, moving either forward from 0 to 12, or backwards from 12 to 0. 3 jumps of 4 3 jumps of 4 or 3. Count the number of jumps made, which tells how many groups of 4 are in 12. There are 3 jumps of 4 to make 12. The quotient (answer) is 3. $12 \div 4 = 3$ Example #2 Solve: 18 ÷ 3 Steps: 1. Draw a number line and label it with marks for every number from 0 to the total of 18. 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 2 Ś 2. Draw as many jumps of 3 as possible moving either forward from 0 to 18, or backwards from 18 to 0.



• This model is difficult to use with a story problem where you want to make a given number of groups, and need to find out how many are in each group.

# Using Number Lines to Divide

This strategy is used with students who are first learning to divide. Students can draw number lines and represent the equal groups or sets as equal jumps along the line. This strategy provides students with a visual model of division.



33

# **Arrays for Division**

Because multiplication and division are opposite operations, the array model used for multiplication can also be used to divide. Arrays are rectangular arrangements of symbols, shapes or objects in equal rows and columns. In an array, the number of rows × the number of columns = the total objects. Students can create arrays to determine the number of groups or number in each group within a total.

<b>Example #1:</b> Solve: 28 ÷ 4 In Context: Angela is putting photos into a photo album. She has 28 photos, and the sheets in the album each hold 4 photos. How many sheets will she fill?				
Steps:				
Draw an array with equal rows of 4 to see how many groups of 4 it takes to make 28.				
<ol> <li>Draw 4 rectangles in the 1st row to stand for the 4 photos on she</li> <li>Draw 4 rectangles in the 2nd row to stand for the 4 photos on she</li> <li>Draw 4 rectangles in the 3rd row to stand for the 4 photos on she</li> </ol>	eet 2. $\rightarrow$ $\bigcirc$			
4. Keep drawing rows of 4 until you reach the total of 28 photos. $\longrightarrow$ $\square$ $\square$ $\square$ $28$				
5. Count how many rows of 4 it took to make 28.				
Answer: 7 rows of 4 make 28. 28 ÷ 4 = 7 A	ngela will fill 7 pages.			
<b>Example #2: Solve:</b> 15 ÷ 3 <b>In Context:</b> Three sisters work together to earn some money raking leaves. They earned \$15. How many dollars go to each sister?				
<b>Steps:</b> Draw an array with 3 equal rows to see how many dollars each sister will get (in each row).				
<ol> <li>Draw 1 rectangle (dollar bill) in each of the 3 rows to give each sister \$1.</li> </ol>	Sister #1 Sister #2 Sister #3			
2. Add a 2nd rectangle in each row to give each sister another \$1.	Sister #1 Sister #2 Sister #3 Sister #3			
3. Add a 3rd rectangle in each row to give each sister another \$1.	Sister #1 Sister #2 Sister #3			
<ol> <li>Keep giving each sister a dollar until all \$15 have been handed out.</li> </ol>	Sister #1   Image: Constraint of the second secon			
5. Count how many rectangles (dollars) are in each row.				
	ter will get \$5.			
Grade 3				

# **Arrays for Division** (Continued with Remainders)



Some problems have numbers that may not divide evenly, such as 11 divided by 5. This situation leaves you with a partial group of extra items. In an array, extra items leave you with an unfinished row or column. The number of leftover items that cannot form a complete group or cannot be shared with each existing group is called the **remainder**.

### **Example #3: Solve:** 36 ÷ 7

In Context: Ronnie's birthday party is in 36 days. He wants to know how many weeks until his birthday, so his dad tells him to divide 36 by 7 days in each week. How many weeks make 36 days?

### Steps:

Draw an array with equal rows of 7 to see how many sevens it takes to make a total of 36.

- 1. Draw 7 circles in the 1st row to stand for the 7 days in the 1st week.  $\rightarrow OOOOOOO7$
- 2. Draw 7 circles in the 2nd row to stand for the 7 days in the 2nd week. → OOOOOOO <sup>14</sup>
- 3. Keep drawing rows of 7 days until you reach the total of 36 days.
- 000000 21
- 000000 28 000000 35

→ O 36

5 rows make 35 days, so you need 1 extra day to make 36 days. -

Count how many rows of 7 it took to make 35.

There are 5 rows with 1 extra circle. This means there is a remainder of 1.

Answer: There are 5 rows (weeks) and a remainder of 1 extra day.

 $36 \div 7 = 5$  with a remainder of 1. Ronnie's party will be in 5 weeks and 1 day.

Tips:

- It is great to use everyday objects to form arrays. Divide up everyday household items into equal rows and columns to answer simple division questions.
- Students can create arrays with simple dots, circles, boxes, or x's to represent anything. •
- Graph paper is a terrific tool for modeling larger problems such as 96 divided into 6 equal rows.
- Remember that you can orient a rectangular array either vertically or horizontally. •
- This is not an efficient model for long term use. It is a learning tool for students beginning to divide.

Grade 4

# **Partial Quotients**


Partial quotients let students divide the total in manageable chunks. Students can multiply by larger numbers and solve the problem quickly in a few steps, or take more steps using smaller numbers, based on their confidence and proficiency level.



### **Solve:** 152 ÷ 4

**In Context:** Parents made 152 cookies for a bake sale at school. They are to be packed into bags that are each filled with 4 cookies. How many bags of cookies will be filled for the parents to sell?

In each step using partial quotients, multiply to make as many groups of 4 as possible, without going over 152. After multiplying, subtract the amount of cookies already grouped into bags. Keep making groups and subtracting them until you run out of cookies. Then add the number of groups (bags) of cookies you have made to find the quotient (ans.



Tip: There is always more than one path to the solution, and any accurate path is acceptable. Turn to the following page to see the above problem solved 3 more ways using partial quotients.


Grades 4-5

# **Partial Quotients (Continued)**

**Solve:** 152 ÷ 4

**In Context:** Parents made 152 cookies for a bake sale at school. They are to be packed into bags that are each filled with 4 cookies. How many bags of cookies will be filled for the parents to sell?





### Tips:

- Encourage students who may be only multiplying by 1, 2, or 10 to begin trying other factors as they are ready.
- As students begin to understand how to use multiplication and division facts and estimation to their advantage, this method can become very efficient, as seen in the last example above.
- Long division is not introduced until 6th grade.

Grades 4-5

# **Multiplying Up**

Multiplication and division are opposite operations so students can practice estimation and use multiplication skills to solve division problems with the "multiplying up" strategy. Here groups are multiplied together, adding more and more groups until the total is reached.




### Solve: 400 ÷ 16

**In Context:** The public library received a donation of \$400 for new books. They want to order as many books as they can. If the books are \$16 each, how many books could they order?

**Steps:** To multiply up, ask yourself, "How many groups of 16 dollars can I make, that will get me as close to \$400 as possible without going over?" Continue making more groups of \$16 until we reach \$400.

First, we will multiply to make groups of 16 dollars, trying get to 400 total. It helps to imagine you have already spent that money on books. Each time we multiply to make the groups of 16 dollars to buy books, we will add that amount to keep track of how much we have spent.




# **Multiplying Up (Continued)**

**Solve:** 400 ÷ 16

**In Context:** The public library received a donation of \$400 for new books. They want to order as many books as they can. If the books are \$16 each, how many books could they order?




Here is the same problem,  $400 \div 16$ , solved four more ways:



Tips:

- Encourage students who may be multiplying by only 1, 2, or 10 to begin trying other factors as they are ready.
- Remember that this approach is not about solving efficiently at first. It is about deepening students' understanding and mathematical flexibility.
- As students begin to understand how to use multiplication and division facts, as well as estimation, this method can become much more efficient, as seen in the bottom middle example above.

Grades 4-5



Grades 4-5

40